

# Temperature dependence of electrical conductance in 1D electron system formed on liquid helium surface

Hideki Yayama <sup>a,1</sup>, Igor B. Berkutov <sup>b</sup>

<sup>a</sup> Department of Physics, Faculty of Science, Kyushu University, 4-2-1 Ropponmatsu, Fukuoka 810-8560, Japan

<sup>b</sup> Institute for Low Temperature Physics and Engineering, Academy of Sciences of Ukraine, 47 Lenin Avenue, 61164 Kharkov, Ukraine

---

## Abstract

The electrical conductance of 1D electrons confined in the channels created on the surface of liquid helium was measured as a function of temperature down to 50 mK. A peak around 0.2 K, which was the evidence for 1D character originating in the intersubband transitions, was observed for the first time. The result of the temperature dependence is in agreement with an existing theory for 1D electron system. Below 0.2 K, the electrons occupy the ground state and behave as a nearly ideal 1D system.

*Key words:* 1D electrons; conductivity; surface electrons; superfluid helium

---

The quasi-one-dimensional (Q1D) electron system has been created by confining electrons in the linear channels formed on the curved helium surface [1–4]. This system has an advantage that the impurities and defects are free in comparison with the solid state one. Sokolov *et al.* [7] calculated the mobility for this system as a function of temperature. According to their theory, the characteristic point for this system is that there exist a maximum in the mobility around 0.2 K. However, this behavior has not been observed yet.

The real experiments are different from the ideal theory. The systems created so far are still narrow 2D because the width of the channel is larger than the inter-electron distance. The electrons near the edge of the channel contribute to the conductance. In addition, since the temperature is high, the system is not in the ground state and it is not strictly a pure 1D.

It was found that the electrons near the edge of the channel localize at low temperatures [5]. Therefore, it is expected that the electrons only in the center are mobile and the 1D character can be observed, if the temperature is lowered below a few hundreds mK.

In this paper, we show the experimental results measured down to 50 mK. We observed a peak at 0.2 K in the conductance for the first time, which originates in the intersubband transitions. This is the clear evidence for the pure 1D electron system in the ground state.

The important part of the experimental cell is a glass optical-diffraction-grating that is used to create the 1D channels. It has 2000 parallel linear grooves on the surface. The cross-section view of the groove shows a very shallow V-shape. The distance between the adjacent grooves is 5  $\mu$ m, the width of the groove is 3.8  $\mu$ m, and the depth is 0.2  $\mu$ m at the center. The superfluid helium comes up to the surface of the grating under the action of the capillary force and fills the grooves. The helium surface sags 0.05  $\mu$ m at the center of the groove by a balance of the surface tension and the gravity.

The grating was placed on the lower electrodes which was DC-grounded. A negative DC voltage was applied to the upper electrode positioned several mm apart from the grating to generate a vertical electric field. The lower electrode was divided into three segments to measure the electron's conductance. An AC voltage was applied to the left segment to drive electrons laterally along the channels and the response was detected

---

<sup>1</sup> Corresponding author. E-mail: yayama@rc.kyushu-u.ac.jp

by the right segment coupled to the electrons capacitively. The central segment was grounded to minimize the cross talk. The conductance of the electrons was calculated from the measured amplitude and the phase shift.

Different from the 2D case, the 1D electrons' motion perpendicular to the channel is quantized and the discrete subbands are formed, because they are subjected to the parabolic confinement potential. According to the theoretical calculation [7], below 0.2 K the electrons are in the ground state and the conductance increases as the temperature increases. But above 0.2 K the conductance decreases with increasing temperature when both the intersubband transitions and the increase of the population of electrons in higher subbands become dominant. As a result, the conductance shows a maximum near 0.2 K. If the maximum is found, it is a strong evidence for the 1D character.

The 2D conductance is known to be a monotonic increasing function as the temperature is lowered [6]. However, our obtained result is quite different from that of 2D electrons. Lowering the temperature from 1.8 K, the measured conductance increases first, like the 2D case, but begins to decrease through a maximum at 1.4 K. The decrease becomes gentle from 0.7 K down to 0.2 K. Below 0.2 K, it starts decreasing again.

To explain this complicated behavior, we consider the contribution of electrons existing near the edge of the grooves [8]. In our experimental condition, the width of the grating is larger than the inter-electron distance and so a few electrons distribute over the width of the groove. In this sense, the system is not pure 1D but Q1D. But the electrons in the center of the groove are expected to behave as pure 1D electrons, because the helium thickness is large and hence the image force acting from the glass grating is small. In contrast, the electrons near the edge of the groove are captured by the random image force caused from the surface roughness of the grating because the helium thickness near the edge is small. For this reason, the electrons near the edge localize due to the potential irregularities at low temperatures.

Based on the experimental facts [5], we can express the measured conductance as a sum of the conductance of the pure 1D electrons,  $1/R_{1D}$ , in the center of the groove and that of the 2D electrons near the edge of the groove,  $1/(R_L + R_G)$ . Here,  $R_L$  is the resistance caused by the localization due to the potential irregularities and  $R_G$  is the resistance caused by the collision with the helium gas. The effect of electron-ripllon scattering is very small compared with the localization effect and it is neglected. These resistances are expressed as  $R_L = R_{L0}T \exp(4.2/T)$  [5] and  $R_G = R_{G0} \exp(-8.5/T)$  [6], where the numerical values are determined from the data in each experiment,  $R_{L0}$  and  $R_{G0}$  are adjustable parameters and  $T(K)$  is the absolute temperature. The




Fig. 1. Temperature dependence of the conductance. The vertical electric field is  $E_\perp = 1000$  V/cm. Dots are experimental values and the solid line is a calculation [7].

conductance of the pure 1D electrons,  $1/R_{1D}$ , is obtained by subtracting  $1/(R_L + R_G)$  from the measured conductance.

In Fig. 1, the experimental results of the conductance of the pure 1D electrons are plotted with dots as a function of temperature. The theoretical conductance calculated from the Sokolov *et al.*'s mobility data [7] is also depicted by a solid line. A factor is adjusted to agree with the experimental data at  $T = 0.2$  K. It appears that the experimental data essentially agree with the theoretical temperature dependence. Above 0.8 K, the conductance is governed with the scattering by helium gas atoms. Below 0.8 K, the electron-ripllon scattering is dominant. The most striking feature is the existence of a maximum at 0.2 K in agreement with the theory. This is the direct evidence for the 1D character which is observed for the first time.

Below 0.2 K, the electrons are in the ground state and therefore it is the ideal pure 1D electron system.

## References

- [1] Yu.Z. Kovdrya, V.A. Nikolaenko, Sov. J. Low Temp. Phys. **18** (1992) 894.
- [2] H. Yayama, A. Tomokiyo, Czech. J. Phys. **46** (1996) 353.
- [3] A.M.C. Valkering, P.K.H. Sommerfeld, P.J. Richardson, R.W. van der Heijden, A.T.A.M de Waele, Czech. J. Phys. **46** (1996) 321.
- [4] R.J.F. van Haren, G. Acres, P. Fozooni, A. Kristensen, M.J. Lea, P.J. Richardson, A.M.C. Valkering, R.W. van der Heijden, Physica **B249-251** (1998) 656.
- [5] H. Yayama, A. Tomokiyo, O.I. Kirichek, I.B. Berkutov, Yu.Z. Kovdrya, Low Temp. Phys. **23** (1997) 878.
- [6] Y. Iye, J. Low Temp. Phys. **40** (1980) 441.
- [7] S.S. Sokolov, G.-Q. Hai, N. Studart, Phys. Rev. B **51** (1995) 5977.
- [8] H. Yayama, I.B. Berkutov, A. Tomokiyo, Physica **B284-288** (2000) 1914.