

Vortex Nucleation and texture of rotating $^3\text{He-A}$ in cylindrical cells with $R \simeq 10\xi_D$.

R. Ishiguro^{a,1}, M. Yamashita^a, T. Igarashi^b, E. Hayata^a, O. Ishikawa^c, Y. Sasaki^a, K. Fukuda^e, M. Kubota^b, H. Ishimoto^b, T. Mizusaki^a, T. Ohmi^a, T. Takagi^d, R. E. Packard^f

^aDepartment of Physics, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan

^bInstitute for Solid State Physics, University of Tokyo, Chiba 277-8581, Japan

^cGraduate School of Science, Osaka City University, Osaka 558-8585, Japan

^dDepartment of Applied Physics, Fukui University, Fukui 910-8507, Japan

^eCollege of Medical Technology, Kyoto University, Kyoto 606-8507, Japan

^fDepartment of Physics, University of California, Berkeley, CA94720, USA

Abstract

We measured cw-NMR at 700 kHz for rotating $^3\text{He-A}$ at 31.5 bar confined in two sizes of cylinders with 0.1 and 0.2 mm diameters at rotation speeds up to 6.28 rad/sec. The NMR signals varied for the different size cylinders. Nucleation of the single vortex was observed in the 0.2 mm cylinder above a critical rotational speed and hysteresis appeared for vortex nucleation and annihilation when the rotational speed was changed. On the other hand in 0.1 mm cylinder, a vortex did not appear up to 6.28 rad/sec, and the NMR absorption gradually changed as the rotation speed increased.

Key words: Superfluid $^3\text{He-A}$; NMR; Texture; Vortex

We studied rotationg superfluid $^3\text{He-A}$ in cylindri-
cal sample cells under rotation. This work is an exten-
sion of the previous experiment[1] but was carried out
under higher rotation speed up to 6.28 rad/sec in sub
mK temperature range, and by using the improved cw-
NMR.

We prepared two cylindrical cells with 0.1 mm and
0.2 mm diameters which were about 10 times larger
than the dipole coherence length $\xi_D \sim 10 \mu\text{m}$ and about
150 cylindrical samples were put together in each cell
and their cylinder axis was parallel to the rotation axis
and also parallel to the magnetic field.

We observed (1) three kinds of NMR satellite signals
in the 0.1 mm diameter sample cell which came from
localized spin waves in textures and (2) one kind of

textural satellite signal and a doubly-quantized non-
singular vortex signal in 0.2 mm diameter sample cell.
There was no vortex nucleation observed in the 0.1
mm diameter cell up to 6.28 rad/sec. In this report we
present the results on nucleation and annihilation of a
single vortex in the 0.2 mm diameter sample cell.

Fig. 1 shows a typical NMR absorption spectrum
observed for 0.2mm sample at $T/T_c = 0.75$ where the
dotted line is the cw-NMR adsorption spectrum at rest
and the solid one is the spectrum at 6.28 rad/sec. The
spectrum is characterized by a parameter R_t^2 defined
by $\Delta f = R_t^2 (f_L^A)^2 / 2f_0$, where Δf is the frequency shift
from the Larmor frequency f_0 and f_L^A is the tempera-
ture dependent longitudinal frequency in the A phase.

The characteristic feature of the spectrum at rest
is that there is a peak at $R_t^2 = 1$ and the absorption
spectrum continuously extended toward f_0 due to the
texture. The solid line taken under 6.28 rad/sec has

¹ Corresponding author. E-mail: ishiguro@scphys.kyoto-u.ac.jp




Fig. 1. NMR absorption line shapes for the 0.2mm sample at $T/T_c = 0.75$ as function of frequency shift from Lari frequency. The dotted line was taken at rest and the solid line was taken at rotation speed of 6.28 rad/sec .

2 peaks. The main peak is located at $R_t^2 = 1$ and a satellite peak is located at $R_t^2 \sim 0.37$. The main peak comes from the texture, which is the same as the second line. The satellite peak appeared at a certain critical rotation speed when we increased the speed and it disappeared at a different critical rotation speed when we reduced the speed from the vortex state. The value $R_t^2 \sim 0.37$ is very close to the doubly quantized non-singular vortex which are observed in bulk $^3\text{He-A}$ [2].

Fig. 2 shows the change in NMR absorption signal as a function of the rotation speed Ω . Since the main peak height is most sensitive to the change of the texture, the main peak height is plotted against rotation speed. Solid circles were taken as the speed increased and the crosses when the speed decreased. Data in Fig. 2 (a) were taken when the sample was cooled to superfluid state at rest and data in (b) when the sample was cooled to the superfluid state under a rotation speed of $+2\text{ rad/sec}$. In Fig. 2 (a), data of solid circles change rapidly around $\Omega_{c1} = 5.0\text{ rad/sec}$ and data of crosses changes rapidly at $\Omega_{c2} (\sim 3.5\text{ rad/sec})$. Data are rather symmetric for both directions of rotation. It was found that the sudden change in the main peak height corresponded to sudden change in the satellite absorption and was caused by the textural change due to introduction of a single vortex in the cell. In Fig. 2 (b), data for a positive rotation are not symmetric against negative rotation. This asymmetry can be attributed to the textural memory effect [3] at the onset of superfluidity.

This nucleation and annihilation phenomena can be interpreted as the same mechanism applied for the ^4He [4] case. The edge of our cylindrical cells were connected with the bulk liquid, and when there were no vortex in the 0.2 mm sample cell, there were many vortices in the bulk. For this reason, in order to nucleate the vortex in 0.2 mm cell, it is not needed to exceed a critical velocity v_c which makes the texture unstable. The nucleation and annihilation of the vor-




Fig. 2. NMR absorption at $R_t^2 = 1$ during acceleration (solid circle) and deceleration (cross). (a): the sample was cooled to superfluid at rest and (b): it was cooled under the rotation speed of $+2\text{ rad/sec}$

tex in a cylindrical cell are determined as follows: The free energy $E'(r) = E(r) - \Omega L$ under a rotation Ω can be calculated as a function of the position of the vortex r from the center of the cylindrical cell. The vortex nucleation occurs when the free energy $E'(r)$ at the center $r = 0$ becomes negative above a certain rotation speed Ω_{c1} . The critical angular velocity Ω_{c1} is calculated by vortex line approximation, and is given by $\Omega_{c1} = n\kappa/(2\pi R^2) \ln(R/a)$, where n is the quantum number of circulation 2, κ is the circulation, R is the radius of the cylinder and a is roughly the vortex core radius $\simeq \xi_D (10\text{ }\mu\text{m})$. For 0.2 mm cell, Ω_{c1} becomes 4.8 rad/sec . When the rotation speed is above Ω_{c2} , $E'(r)$ is the minimum at $r = 0$ and the maximum in $E'(r)$ exists somewhere between $r = 0$ and R . When rotation speed is below $\Omega_{c2} = n\kappa/(2\pi R^2)$, the energy barrier disappears. For 0.2 mm cell, Ω_{c2} becomes 2.0 rad/sec . Therefore, the nucleation of the vortex is caused by introduction of the pre-existing vortices in bulk sample into the cylindrical cell and the annihilation of the vortex is due to disappearance of the energy barrier in $E'(r)$ at Ω_{c2} .

Agreement of the calculation with experiment is good. The texture and the large core size of the texture should be considered in the calculation to explain the textural memory effect and to get better agreement with experiment.

This joint research was carried out at ISSP, University of Tokyo.

References

- [1] O. Ishikawa, *et al.*, Physica B **284-288** (2000) 248.
- [2] O. V. Lounasmaa, E. Thuneberg, Proc. Natl. Acad. Sci. USA **96** (1999) 7760.
- [3] A. J. Manninen, *et al.*, Phys. Rev. Lett. **77** (1996) 5086.
- [4] R. E. Packard, *et al.*, Phys. Rev. A. **6** (1972) 799.